

SYMBOLIC COMPUTATION OF THE LEGENDRE–FENCHEL CONJUGATE AND BICONJUGATE OF A QUARTIC UNIVARIATE POLYNOMIAL

YVES LUCET

ABSTRACT. Given a quartic univariate polynomial P , we compute symbolically its Legendre–Fenchel biconjugate and conjugate.

The Legendre–Fenchel conjugate of a univariate function P is defined as

$$(1) \quad P^*(s) := \sup_{x \in \mathbb{R}} [sx - P(x)].$$

We want to compute *explicitly* the point $\hat{X}(s)$ where this maximum is attained when P is a quartic univariate polynomial.

The key idea is to note that the biconjugate or convex hull P^{**} of P is obtained by adding a bitangent to $\text{Epi}P$ (see Figure 1). In other words,

$$(2) \quad P^{**}(x) = \begin{cases} P(x) & \text{if } x \notin [x_1, x_2], \\ P'(x)(x - x_1) + P(x_1) & \text{otherwise;} \end{cases}$$

with $x_1 \neq x_2$ satisfying $P'(x_1) = P'(x_2)$ and $x_1P'(x_1) - P(x_1) = x_2P'(x_2) - P(x_2)$ (see Figure 1).




FIGURE 1. Computing the biconjugate of P amounts to adding bitangents to $\text{Graph}P$

Date: September 30, 1998.




FIGURE 2. The biconjugate and the Legendre–Transform of a quartic polynomial: $P(x) = x^4 - 7x^3 - 40x^2 + 76x + 240$

In addition, we can parametrize $\text{Graph}P^*$ by

$$(3) \quad \begin{cases} s &= P'(x) \\ P^*(s) &= sx - P(x) \end{cases} \quad x \in (-\infty, x_1] \cup [x_2, +\infty).$$

So computing P^{**} amounts to finding x_1 , x_2 , and $\bar{u} := P'(x_1)$.

If we let the parameter x describe the whole real line in (3), we obtain a contact transformation (see Figure 2). To obtain P^* , we must remove a part of the graph of the function $F : x \mapsto (P'(x), xP'(x) - P(x))$. We note that if $\alpha \leq \beta$ are the real roots of P'' (if there is no real root, P is convex and we are done), the “singular” points of $\text{Graph}F$ are $F(\alpha)$ and $F(\beta)$.

All in all, computing the convex hull amounts to computing x_1 and x_2 . Next, solving a 3rd degree polynomial equation gives the usual nonparametric representation of the conjugate.

1. COMPUTING THE CONVEX HULL

We define

$$\begin{aligned} P_1(x) &:= u - P'(x), \\ P_2(x) &:= v - xP'(x) + P(x). \end{aligned}$$

The line $y = ux - v$ is tangent to $\text{Graph}P$ at x (see Figure 1). We are looking for $x_1 \neq x_2$ such that $F(x_1) = F(x_2)$. In other words, x_1 and x_2 are solutions of the polynomial system

$$\begin{cases} P_1(x) = 0, \\ P_2(x) = 0. \end{cases}$$

We deduce that the resultant $Q(u, v)$ of P_1 and P_2 is null (the function $Q(u, v)$ is null if and only if P_1 and P_2 have a common root). It allows us to eliminate x between both equations to obtain an implicit representation of $\text{Graph}F$: $Q(u, v) = 0$.

The next step is to note that if $\nabla Q(\bar{u}, \bar{v}) \neq 0$, there is no multiple point in a neighborhood of (\bar{u}, \bar{v}) . Indeed, from the Implicit Function Theorem $Q(u, v) = 0$

can be locally written as $v = q(u)$ with q a smooth function. So there can be no double point in a neighborhood of (\tilde{u}, \tilde{v}) .

Lemma. *All multiple point of Graph F lie in*
 $S := \{(u, v) \in \mathbb{R} \times \mathbb{R} : \nabla Q(u, v) = 0 \text{ and } Q(u, v) = 0\}$.

Finally S contains $F(\alpha)$ and $F(\beta)$ and can have at most 3 points. Hence the remaining one is (\bar{u}, \bar{v}) defining the bitangent.

Eventually

- First, compute $Q(u, v)$ the resolvent of P_1 and P_2 .
- Next, solve $\nabla Q(u, v) = 0$ and deduce (\bar{u}, \bar{v}) .
- Finally, solve $P'(x) = \bar{u}$ to obtain x_1 and x_2 .

If we apply our method to a quartic polynomial, we deduce the following result.

Proposition (Convex hull computation). *Let $P(x) := x^4 + a_3x^3 + a_2x^2 + a_1x + a_0$. If $3a_3^2 - 8a_2 \leq 0$, P is convex: $P^{**} = P$.*

Otherwise $3a_3^2 - 8a_2 > 0$ and

$$\begin{aligned}\bar{u} &= -\frac{1}{2}a_3a_2 + \frac{1}{8}a_3^3 + a_1 \\ \bar{v} &= \frac{1}{64}a_3^4 - \frac{1}{8}a_2a_3^2 + \frac{1}{4}a_2^2 - a_0 \\ x_1 &= -\frac{1}{4}a_3 - \frac{1}{4}\sqrt{3a_3^2 - 8a_2} \\ x_2 &= -\frac{1}{4}a_3 + \frac{1}{4}\sqrt{3a_3^2 - 8a_2}\end{aligned}$$

Proof. Only the last part is non-trivial. To prove it, check directly that equalities $P'(x_1) = P'(x_2)$ and $x_1P'(x_1) - P(x_1) = x_2P'(x_2) - P(x_2)$ hold with $x_1 \neq x_2$ (in fact, $x_1 < \alpha < \beta < x_2$, where $\alpha < \beta$ are the real roots of $P''(x) = 0$). \square

2. COMPUTING THE CONJUGATE

As for now, we only know a parametric representation of P^* , namely (3). Obtaining an explicit writing implies solving a 3rd degree polynomial equation.

We summarize a well-known method to do it. Using $X := x + a_3/4$, we transform the equation $P'(x) = s$ into

$$(4) \quad X^3 + pX + q - \frac{s}{4} = 0,$$

with $p := \frac{1}{4}(2a_2 - \frac{3}{4}a_3^2)$ and $q := \frac{1}{4}(\frac{1}{8}a_3^3 - \frac{1}{2}a_2a_3 + a_1)$. Now, denoting $j := \exp(2i\pi/3)$ the (usual) cubic root of 1, we have

$$(X - A - B)(X - jA - j^2B)(X - j^2A - jB) = X^3 - 3ABX - (A^3 + B^3).$$

Hence we are looking for $X := A + B$ with $AB = -p/3$ and $A^3 + B^3 = s/4 - q$. Thus A^3 and B^3 are roots of the resolvent equation: $X^2 + (q - s/4)X - p^3/27 = 0$.

Let $\Delta := (q - s/4)^2 + 4p^3/27$ and consider the following three cases.

Case $\Delta > 0$ There are two distinct roots of the resolvent equation. The unique real root of (4) is given by Cardan's Formula:

$$(5) \quad \hat{X}(s) := \left[\frac{1}{2} \left(\frac{s}{4} - q + \sqrt{\left(q - \frac{s}{4} \right)^2 + \frac{4}{27} p^3} \right) \right]^{1/3} + \left[\frac{1}{2} \left(\frac{s}{4} - q - \sqrt{\left(q - \frac{s}{4} \right)^2 + \frac{4}{27} p^3} \right) \right]^{1/3}$$

Case $\Delta = 0$ There is a double root of the resolvent equation. Equation (4) has two distinct real roots. The unique *single* real root is

$$(6) \quad \hat{X}(s) := 2 \left(\frac{1}{2} \left(\frac{s}{4} - q \right) \right)^{1/3}.$$

A continuity argument implies that the single root is the one we are looking for.

Case $\Delta < 0$ Define $r := 1/2(s/4 - q + i\sqrt{-\Delta})$ with i the usual square root of -1 . Name γ a cubic root of r . Equation (4) has three distinct real roots: $\gamma + \bar{\gamma}$, $j\gamma + \overline{j\gamma}$, and $j^2\gamma + \overline{j^2\gamma}$ (where overline denotes complex conjugacy). In fact, Galloï's theory tells us that one cannot write these real roots with only *real* radicals¹.

The root we are looking for is given by the Maple function `surd`:

$$(7) \quad \hat{X}(s) := \text{surd} \left(\frac{1}{2} \left[\frac{s}{4} - q + i \sqrt{-\left(q - \frac{s}{4} \right)^2 - 4 \frac{p^3}{27}} \right], 3 \right) + \text{surd} \left(\frac{1}{2} \left[\frac{s}{4} - q + i \sqrt{-\left(q - \frac{s}{4} \right)^2 - 4 \frac{p^3}{27}} \right], 3 \right).$$

Indeed, the same continuity argument as above implies that we want the cubic root of r *closest* to the real axis. This is precisely the root the function `surd` computes.

Hence Formulas (5)–(7) allow us to compute the conjugate of P .

Theorem (Conjugate computation). *Let $P(x) := x^4 + a_3x^3 + a_2x^2 + a_1x + a_0$. If $3a_3^2 - 8a_2 \leq 0$, P is convex, $\Delta \geq 0$, and the point where the maximum is attained in (1) is given by Formulas (5) and (6).*

Otherwise $3a_3^2 - 8a_2 > 0$, the convex hull of P is given by Formula (2) with x_1 and x_2 as in the previous proposition. The point where the maximum is attained in (1) is given by Formulas (5)–(7).

Proof. Only the convex case needs to be proved. Since $p = 1/16(8a_2 - 3a_3^2)$ is non-negative, $\Delta = (q - s/4)^2 + 4p^3/27$ is non-negative too, and the theorem follows. \square

Example 1. *Take $P(x) := (x^2 - 1)^2 = x^4 - 2x^2 + 1$. We easily compute $\bar{u} = \bar{v} = 0$, $x_1 = -1$, $x_2 = 1$, $p = -1$, and $q = 0$. The resolvent equation is: $X^2 - sX/4 + 1/27 = 0$ and the discriminant $\Delta = -4/27 + 1/16s^2$.*

¹To compute the real and imaginary part of r , one has to solve a 3rd degree equation, thus coming back to the same problem. For that reason that case was referred to as “casus irreducibilis”.




FIGURE 3. Removing two of the three branches of $P'(x) = s$ gives the conjugate

Consequently, if Δ is non-negative the solution is

$$\hat{X}(s) := \left(\frac{1}{8}s + \frac{1}{72}\sqrt{81s^2 - 192} \right)^{1/3} + \left(\frac{1}{8}s - \frac{1}{72}\sqrt{81s^2 - 192} \right)^{1/3},$$

otherwise Δ is negative and the solution is

$$\hat{X}(s) := 2 \operatorname{Re}(\operatorname{surd}(\frac{1}{8}s + \frac{1}{72}I\sqrt{192 - 81s^2}, 3));$$

where Re denotes the real part of a complex number.

CECM, DEPARTMENT OF MATHEMATICS AND STATISTICS,, SIMON FRASER UNIVERSITY, BURN-
 ABY BC, V5A 1S6

E-mail address: lucet@na-net.ornl.gov